

EXHIBIT A

Polymers in Drug Delivery

EDITED BY

Ijeoma F. Uchegbu
Andreas G. Schätzlein

TEMPLE UNIVERSITY
HEALTH SCIENCE CENTER LIBRARY
SOUTH



Taylor & Francis

Taylor & Francis Group
Boca Raton London New York

CRC is an imprint of the Taylor & Francis Group,
an informa business

Cover illustration by Meilan Pang, Simon Fraser University, Vancouver, Canada.

Published in 2006 by
CRC Press
Taylor & Francis Group
6000 Broken Sound Parkway NW, Suite 300
Boca Raton, FL 33487-2742

© 2006 by Taylor & Francis Group, LLC
CRC Press is an imprint of Taylor & Francis Group

No claim to original U.S. Government works
Printed in the United States of America on acid-free paper
10 9 8 7 6 5 4 3 2 1

International Standard Book Number-10: 0-8493-2533-1 (Hardcover)
International Standard Book Number-13: 978-0-8493-2533-5 (Hardcover)
Library of Congress Card Number 2005046684

This book contains information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. A wide variety of references are listed. Reasonable efforts have been made to publish reliable data and information, but the author and the publisher cannot assume responsibility for the validity of all materials or for the consequences of their use.

No part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access www.copyright.com (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC) 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

Trademark Notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Library of Congress Cataloging-in-Publication Data

Polymers in drug delivery / edited by Ijeoma Uchegbu, Andreas Schatzlein.
p. cm.

Includes bibliographical references and index.

ISBN 0-8493-2533-1 (alk. paper)

1. Polymeric drug delivery systems. 2. Polymeric drugs. 3. Polymers. 4. Drug delivery systems. I. Uchegbu, Ijeoma F. II. Schatzlein, Andreas.

RS201.P65P6444 2006
615'.3--dc22

2005046684

informa

Taylor & Francis Group
Is the Academic Division of Informa plc.

Visit the Taylor & Francis Web site at
<http://www.taylorandfrancis.com>
and the CRC Press Web site at
<http://www.crcpress.com>

Prefac

H. Ringsdorf and

Drug Delivery

Not only from a evolves and grow part of a paradig moved from syn macologically ac materials science

When in 195 field of macromo at that time hard; Lipmann reporter ably even more li rang in molecular in the 1920s, the 1 industry and incl daily use. The "P. new materials. T ment, synthetic fi venously (i.v.) inj widely on both si

The possibili 1950s with the r biology, protom. molecular recogr and the evolution in many director fathers of polym Staudinger, reflec Today — Aging synthetic cells as of polymer chem beyond which on may give rise to originated from polymer chemist already in 1959, Hermann Staudi dreaming about ' manship so succ mankind were to

Table of Contents

Chapter 1	Introduction.....	1
<i>Ijeoma F. Uchegbu</i>		
Chapter 2	Selecting the Right Polymer for Biomaterial Applications.....	7
<i>Allan S. Hoffman</i>		
Chapter 3	Polymer Characterization Techniques.....	23
<i>Wei Wang</i>		
Chapter 4	The Role of Polymers in Solid Oral Dosage Forms.....	35
<i>Richard A. Kendall and Abdul W. Basit</i>		
Chapter 5	Hydrogel Drug Delivery Systems.....	49
<i>Seong Hoon Jeong, Kang Moo Huh, and Kinam Park</i>		
Chapter 6	An Assessment of the Role of Polymers for Drug Delivery in Tissue Engineering.....	63
<i>Patrick J. Ginty, Steven M. Howdle, Felicity R.A.J. Rose, and Kevin M. Shakesheff</i>		
Chapter 7	Poly(lactic-co-glycolic acid) Microspheres.....	81
<i>Camille Bouissou and Chris van der Walle</i>		
Chapter 8	Polymeric Nanoparticles as Drug Carriers.....	101
<i>Hervé Hillaireau and Patrick Couvreur</i>		
Chapter 9	Polymeric Micelles as Pharmaceutical Carriers.....	111
<i>Vladimir P. Torchilin</i>		
Chapter 10	Polymeric Vesicles.....	131
<i>Ijeoma Florence Uchegbu, Shona Anderson, and Anthony Brownlie</i>		
Chapter 11	Polymer-Drug Conjugates.....	155
<i>Vaikunth Cuchelkar and Jindřich Kopeček</i>		

Chapter 12 Polymers Used for the Delivery of Genes in Gene Therapy	183
<i>Pet Lee Kan, Andreas G. Schätzlein, and Ijeoma F. Uchegbu</i>	
Chapter 13 Dendrimers in Drug and Gene Delivery	199
<i>Christine Dufès, Ijeoma F. Uchegbu, and Andreas G. Schätzlein</i>	
Glossary	237
Index	245

111

lji

CONTENTS

1.1 Drug Delivery
1.2 Polymers
1.3 This Volume
1.4 The Future
References

1.1 DRUG DELIVERY

The science of principles to control drug delivery. When drugs are released at receptors or sites, the "wrong" tissue is targeted. Scientists rescue drug activity and (2)

Drug delivery (1) the emergence of materials with properties that realize that it is sites where they index. Today's work and, preferably, dynamic and be

1.2 POLYMERS

Whether the control of specific cells is an entity. In the "cousins," dendritic systems. Scientists

- Prolong
- Shift drug
- Shunt drug

TABLE 4.4

pH-Sensitive Polymers Commonly Used in the Production of Delayed-Release Oral Dosage Forms

Polymer	Dissolution Threshold pH	Aqueous Dispersion
Cellulose Derivatives		
Cellulose acetate trimellitate	5.0	—
Hydroxypropyl methylcellulose 55	5.5	—
Hydroxypropyl methylcellulose acetate succinate L	5.5	Acoat AS-L
Hydroxypropyl methylcellulose acetate succinate M	6.0	Acoat AS-M
Cellulose acetate phthalate	6.0	Aquacoat CPD
Hydroxypropyl methylcellulose acetate succinate H	6.8	Acoat AS-H
Acrylic Derivatives		
Poly(methacrylic acid, ethyl acrylate) 1:1	5.5	Eudragit L30-D55 Eudragit 30D Kollicon MAE30 DP Acryl-cze
Poly(methacrylic acid, methyl methacrylate) 1:1	6.0	—
Poly(methacrylic acid, methyl methacrylate, methyl acrylate) 2.5:6.5:1	6.8	Eudragit FS
Poly(methacrylic acid, methyl methacrylate) 1:2	7.0	—
Polyvinyl Derivatives		
Polyvinyl acetate phthalate	5.0	Sureteric

Note: All polymers are available in powder/granule form for use in organic solutions and in some cases ready-to-use aqueous dispersions.

A polymer with a dissolution threshold pH in the range 5 to 6 is considered ideal for use as an enteric coat; this is based on the premise that the pH of the stomach, even in the fed state, will rarely reach this level but will exceed this level in the duodenum, where secretion of bicarbonate neutralizes the acidic chyme leaving the stomach.

There is no single enteric polymer that is applicable for the enteric coating of all drug molecules. The nature of the core material (acidity or basicity, or permeability through different enteric polymer films) may limit the choice of polymer. The pK_a of the coating polymer must also be carefully considered, and the potential for premature release in the stomach (for polymers with low pK_a values) weighed against the requirement for a rapid release in the small intestine. Because the physicochemical properties of the drug will have a bearing on this, it is important to consider the consequences of premature release in the stomach (drug degradation or risk of mucosal damage) alongside the requirement for a rapid release of a poorly soluble drug in the small intestine in order to optimise bioavailability and achieve the desired therapeutic effect.

Enteric coating is not without its problems. A lag time of 1.5 to 2 h postgastric emptying for complete disintegration of an enteric-coated capsule and tablet has been demonstrated [6,24]. This is slower than reported for *in vitro* disintegration times, and implies that modified-release dosage forms should be designed as multiple-unit systems, in which the increased surface-area-to-volume ratio would reduce the time for intestinal disintegration while minimizing the possibility of total failure of the dosage form and premature release in the stomach. Furthermore, the *in vivo* evidence highlights the need for new enteric polymers to be developed, which will improve the rapidity